


We claim:

1. A process for the preparation of 2-(4-N,N-dialkylamino-2-hydroxybenzoyl)benzoic esters of the formula I,



15 in which the substituents, independently of one another, have the following meanings:

R¹ and R²

are C₁-C₆-alkyl, C₃-C₁₀-cycloalkyl;


20

R³ is C₁-C₁₂-alkyl, C₃-C₁₀-cycloalkyl

by

- 25 I. reaction of 3-N,N-dialkylaminophenol of the formula II, in which R¹ and R² have the meanings given above, with phthalic anhydride of the formula III to give 2-(4-N,N-dialkylamino-2-hydroxybenzoyl)benzoic acid of the formula IV and

30




- 40 II. subsequent esterification of the 2-(4-N,N-dialkylamino-2-hydroxybenzoyl)benzoic acid of the formula IV formed in stage I with a C₁-C₁₂-alcohol or a cyclic C₃-C₁₀-alcohol in the presence of an acidic catalyst to give the 2-(4-N,N-dialkylamino-2-hydroxybenzoyl)benzoic ester of the formula I,

45

RECEIVED BY
MAY 15 2002

14




which comprises purifying the ester of the formula I formed in a further process stage III by treatment with an adsorbent and/or by distillation.

10

2. A process as claimed in claim 1, wherein the adsorbent is a substance chosen from the group consisting of activated carbons, aluminum oxides, zeolites and silica gels.
3. A process as claimed in claim 1 or 2, wherein the esterification in the process stage II is carried out in the presence of sulfuric acid as catalyst.
4. A process as claimed in any of claims 1 to 3, wherein the 2-(4-N,N-dialkylamino-2-hydroxybenzoyl)benzoic ester of the formula I formed is crystallized prior to the treatment with an adsorbent and/or distillation.
5. A process as claimed in any of claims 1 to 4, wherein the 2-(4-N,N-dialkylamino-2-hydroxybenzoyl)benzoic ester of the formula I formed comprises less than 10 ppm of rhodamine.
6. A process as claimed in any of claims 1 to 5, wherein the benzoic ester is n-hexyl 2-(4-N,N-diethylamino-2-hydroxybenzoyl)benzoate of the formula Ia

35



40

7. A process as claimed in any of claims 1 to 6, wherein, in the process stage III, the adsorbent used is activated carbon or silica gel.
8. A process as claimed in claim 7, wherein, in process stage III, the ester is purified by treatment with activated carbon and subsequent distillation.

45

RECEIVED BY
JAN 24 1967


9. A process as claimed in claim 8, wherein, in the process stage III

- 5 a. the ester is dissolved in a nonpolar solvent at a temperature in the range from 10°C to 100°C,
- b. this solution is passed over a granular activated carbon bed at a temperature in the range from 20°C to 100°C,
- 10 c. the ester, after passing through the granular activated carbon bed, is separated off from the solvent by distillation.

10. A process as claimed in claim 9, wherein the solvent used in the process step IIIa is cyclohexane or toluene.

11. A process for the preparation of n-hexyl 2-(4-N,N-diethylamino-2-hydroxybenzoyl)benzoate of the formula Ia

20




25

Ia

by

- 30 I. reaction of 3-N,N-diethylaminophenol of the formula IIa with phthalic anhydride of the formula III to give 2-(4-N,N-diethylamino-2-hydroxybenzoyl)benzoic acid of the formula IVa,

35




40

- II. esterification of the 2-(4-N,N-diethylamino-2-hydroxybenzoyl)benzoic acid of the formula IVa formed in stage I in hexanol in the presence of sulfuric acid to
- 45 give n-hexyl

REPLACED BY
PAT 34 ANDT

2-(4-N,N-diethylamino-2-hydroxy)benzoyl)benzoate of the formula Ia



10 and isolation of the n-hexyl ester Ia in crystalline form,

III.

- 15 a. dissolution of the n-hexyl ester Ia in toluene or hexanol at a temperature in the range from 25°C to 50°C,
- b. metering of this solution over a granular activated carbon bed or a silica gel bed at a temperature in the
- 20 range from 25°C to 50°C and
- c. subsequent isolation of the n-hexyl ester by separating off the toluene and/or hexanol by distillation.

25

30

35

40

45


REPLACED BY
ACT 34 ANDT

Preparation of 2-(4-N,N-dialkylamino-2-hydroxybenzoyl)benzoic esters

5 Abstract

The invention relates to a process for the preparation of 2-(4-N,N-dialkylamino-2-hydroxybenzoyl)benzoic esters of the formula I,

10




15

in which the substituents R^1 to R^3 , independently of one another, have the meanings given in the description by:

20

- I. reaction of 3-N,N-dialkylaminophenol of the formula II with phthalic anhydride of the formula III to give 2-(4-N,N-dialkylamino-2-hydroxybenzoyl)benzoic acid of the formula IV and


25



30

- II. subsequent esterification of the 2-(4-N,N-dialkylamino-2-hydroxybenzoyl)benzoic acid of the formula IV formed in stage I with a C_1 - C_{12} -alcohol or a cyclic C_3 - C_{10} -alcohol in the presence of an acidic catalyst to give the 2-(4-N,N-dialkylamino-2-hydroxybenzoyl)benzoic ester of the formula I,

35



40

which comprises purifying the ester of the formula I formed in a further process stage III by treatment with an adsorbent and/or by distillation.

45